

Pre- and Postsurgical Detection of IgG, IgM, and IgA Specific to Hydatidosis by ELISA With Purified Antigen Enriched With the 5/B Antigen Complex

Olga Doiz,¹ Rafael Benito,^{1*} Joaquina Gil,¹ Almudena Rojas,² M. Carmen Rubio,¹ and Antonio Osuna³

¹Servicio de Microbiología, Hospital Clínico Universitario, Zaragoza, España

²Vircell SL, Granada, España

³Instituto de Biotecnología, Facultad de Ciencias, Universidad de Granada, Granada, España

An enzyme-linked immunoassay (ELISA) using purified 5/B *Echinococcus* enriched antigen was used to follow IgG, IgM, and IgA antibody levels pre- and posttreatment or surgical removal of hydatid cysts. The sensitivity was 97%, 37.5%, and 54.5%, respectively, and the specificity was 95.7%, 100%, and 98.9%, respectively. All isotypes could be detected 3 years after surgical removal of cysts in patients showing no

remaining cyst evidence. This was especially true for IgG, which persisted in 85.2% of the patients. The data indicate that antigen purification improves specificity without affecting sensitivity, although this new antigen offers no advantages in the postsurgical monitoring of the patients. *J. Clin. Lab. Anal.* 16:295–298, 2002.

© 2002 Wiley-Liss, Inc.

Key words: *Echinococcus granulosus*; hydatidosis; antigens; serological follow-up

INTRODUCTION

The cross-reactions between *Echinococcus granulosus* and other infectious agents are well known and affect the specificity of serological tests. Cross-reactivity has been reported in infections by *E. multilocularis*, *E. vogeli* (1), *Taenia saginata* (2–7), *Taenia solium* (5,8–10), *Hymenolepis nana* (10), *Ascaris lumbricoides* (3), *Trichinella* (11), *Fasciola* (8,11), *Giardia lamblia* (5), *Toxoplasma gondii* (5), *Plasmodium falciparum* (5), and *Leishmania* (4). Furthermore, bacterial infections, such as tuberculosis (8), and host antigens, such as seroalbumin (12,13) and red blood cells (14–16), are sources of false-positive results in hydatidosis serological tests.

Several *Echinococcus* antigen fractions have been tested for improved sensitivity and specificity of serological tests to diagnose hydatidosis (1,10). Preparations enriched with 5 and B fractions were most successful (17,18).

In the present work we used an enzyme-linked immunosorbent assay (ELISA) to study the specific IgG, IgM, and IgA responses to purified hydatid antigen, enriched with the 5/B antigen complex. We analysed the sensitivity and specificity of the 5/B antigen

complex, as well as its possible application in the diagnosis and posttreatment monitoring of human hydatidosis.

MATERIALS AND METHODS

Patients

Group 1

This group was comprised of 72 patients with hydatidosis (36 males and 36 females, 18–82 years old, average 50.1 years, SD = 15.2) who had been diagnosed and monitored in the Hospital Clínico Universitario of Zaragoza between 1994 and 1999. The inclusion criteria were indirect hemagglutination assay (IHA) (Fumouze, Levallois Perret, France) $\geq 1/160$ in at least two sera,

*Correspondence to: Rafael Benito, Servicio de Microbiología, Hospital Clínico Universitario, Avda. San Juan Bosco 15, 50009-Zaragoza, Spain. E-mail: lmic-benito@hcu-iblesa.es

Received 17 January 2001; Accepted 23 July 2002

DOI 10.1002/jcla.10056

Published online in Wiley InterScience (www.interscience.wiley.com).

and the presence of radiologic, ecographic, or tomographic images compatible with hydatidosis.

The cysts were detected in the following locations: 47 liver, five lung, five peritoneum, one spleen, one pancreas, one gluteus, and 12 multiple. In 38 patients, the cysts were not complicated (whole with a crystalline fluid and hyaline aspect); in 28 patients the cysts were complicated (fissured, ruptured, or infected), according to the classification of Verastegui et al. (10); and in six patients the cysts were calcified. During this time, 45 patients underwent surgery, 20 received medical and surgical treatment, two received medical treatment (mebendazole or albendazole), and five patients were not treated. A total of 233 samples were taken, with an average of 4.7 samples per patient (range 2–14). Thirty-three samples (from 33 patients) were taken prior to treatment, and 200 were taken during the monitoring period. During this monitoring period, asymptomatic patients were considered cured after radiologic, ecographic, and tomographic examinations were performed 1 year after treatment.

Group 2

In group 2 we studied cross-reactions in sera from 93 patients: 30 with rheumatoid factor, 10 with antinuclear antibodies, 12 with toxoplasmosis, six with cysticercosis, 12 with leishmaniasis, and 23 with trichinosis.

All samples from both groups were stored in aliquots and kept at -70°C until they were studied.

Techniques

The samples were analysed using commercial ELISA kits (Hydatidosis ELISA IgG, IgM, and IgA; Vircell, SL, Granada, Spain), following the manufacturer's instructions. These kits incorporate a purified antigen enriched with the 5/B antigen complex, according to the methodology of Rogan et al. (19) (modified by Sbihi et al. (17)), and based on the procedures of Oriol et al. (20) using hydatid fluid from sheep-liver cysts.

Briefly, the samples were added to microtitre plates (Nunc Maxisorb, Denmark) containing the antigen, and incubated for 45 min at 37°C . For the IgG determination, 5 μL of the samples were diluted in 100 μL of dilution buffer. To determine the IgA and IgM, 5 μL of the samples were diluted in 75 μL of dilution buffer and 25 μL of anti-human IgG serum. Next, 100 μL of peroxidase-conjugate goat antibody were added to human IgG, IgA, or IgM and incubated for 30 min at 37°C . Finally, 100 μL of tetramethyl-benzidine were added. After incubation for 20 min at room temperature, the reaction was stopped by the addition of 50 μL of sulphuric acid 0.5 M. Absorbance values were measured at 450/620 nm. The results were expressed as

an index (sample absorbance/cut-off). Samples with an index lower than 0.9 were regarded as negative, between 0.9 and 1.1 as a grey-zone result, and higher than 1.1 as positive.

RESULTS

The IgG was positive in 32 of the 33 samples immediately prior to treatment (97%), while the ELISA IgM and IgA were positive in 37.5% and 54.5% of the sera, respectively (Table 1). The only negative sample had an IHA titre of 1/160, a grey-zone result for IgM, and a negative result for IgA. This patient was a 40-year-old female with a 7-cm nonfertile hyaline liver cyst.

The complicated cysts gave the most frequent positive results for all the immunoglobulin isotypes. Sensitivity of IgM and IgA was low in the noncomplicated cysts (37.5% and 43.8%, respectively) but somewhat higher in the complicated ones (50% and 76.9%, respectively). The IgG was positive in all patients with complicated and calcified cysts. No relationship was found between the presence of IgM and the cyst status. In nine of the 65 patients (13.8%) a positive IgM result was detected immediately after the operation, with no significant differences between cured and noncured patients.

In the control group (Table 2), the detection specificity of the different isotypes was 95.7%, 100%, and 98.9% for IgG, IgM, and IgA, respectively. The most frequent sources of positive hydatidosis serology in the absence of *Echinococcus* parasitism were cysticercosis for IgG and leishmaniasis for IgA. It should be noted that leishmaniasis and trichinosis were sources of ELISA grey-zone results.

Of the 65 surgically-treated patients, 38 were considered cured, but in the other 27 the hydatidosis persisted. Cysts persisted in six of the seven patients not surgically treated. One patient was considered cured after cyst calcification and negative serology, 5 years after diagnosis.

In 38 patients who were considered cured, a progressive decline in the number of seropositive cases was observed in all tests, although the rate of negative results varied. The IgA was the test that became negative

TABLE 1. ELISA results in samples of patients with hydatid disease before surgery

Cyst status	No.	EIA IgG Positive/ analyzed	EIA IgM Positive/ analyzed	EIA IgA Positive/ analyzed
Noncomplicated	16	15/16 (93.8%)	6/16 (37.5%)	7/16 (43.8%)
Complicated	13	13/13 (100%)	6/12 (50%)	10/13 (76.9%)
Calcified	4	4/4 (100%)	0/4 (0%)	1/4 (25%)
Total	33	32/33 (97%)	12/32 (37.5%)	18/33 (54.5%)

TABLE 2. Results of hydatid serology in 93 samples of the control group

Patients	IgG		IgM		IgA	
	Positive (%)	GZR (%)	Positive (%)	GZR (%)	Positive (%)	GZR (%)
Rheumatoid factor	0/30 (0%)	0/30 (0%)	ND	ND	0/30 (0%)	1/30 (3.3%)
Antinuclear antibodies	0/10 (0%)	0/10 (0%)	ND	ND	0/10 (0%)	0/10 (0%)
Toxoplasmosis	1/12 (8.3%)	0/12 (0%)	ND	ND	0/12 (0%)	0/12 (0%)
Cysticercosis	1/6 (16.7%)	0/6 (0%)	ND	ND	0/6 (0%)	0/6 (0%)
Leishmaniasis	0/12 (0%)	2/12 (16.7%)	ND	ND	1/23 (4.3%)	0/12 (0%)
Trichinosis	2/23 (8.7%)	0/23 (0%)	0/23 (0%)	1/23 (4.3%)	0/23 (0%)	0/23 (0%)
Total	4/93 (4.3%)	2/93 (2.2%)	0/23 (0%)	1/23 (4.3%)	1/93 (1.1%)	1/93 (1.1%)

GZR, grey zone of reaction; ND, non-determinate.

most rapidly (2.9% positive at 3 years; Table 3). In 27 patients with persistent cysts after treatment, the IgG proved positive in 100% of the cases after 3 years (Table 4), while the IgA and IgM remained positive in around 50% of the cases during the same period. In six patients not surgically treated and with persistent cysts, the IgG remained positive in 100% of the tests over the entire observation period, and the percentages of positive IgA and IgM showed no change (Table 5).

DISCUSSION

The sensitivity of the ELISA-IgG with enriched antigen was equal to or higher than that reported by other authors performing ELISA with crude or lipoprotein antigen, ranging from 87% to 100% (3,5,8,11). Nevertheless, it would also be necessary to study the sensitivity of our ELISA in a group of patients not selected for having other serologically positive tests. The sensitivity of the test with enriched antigen for detecting specific IgM and IgA was insufficient. The values for IgM were lower than those reported by Matossian et al. (11) (58%), Orduña et al. (5) (80.4%), and Candolfi et al. (21) (54%) with crude antigen, but higher than those reported by Tassi et al. (1) with antigen fractions and scolex (with percentages of 27–40% and 33–35%, respectively), according to the antigen fraction considered. In the case of IgA, its values were similar to those found by Force et al. (8) (50%) and Candolfi et al. (21) (53%) using crude antigen.

The specificity of the enzyme-immune assay ranged from 95.7% to 100% according to the isotype of the immunoglobulin considered. Only sera of patients with

TABLE 3. Post-surgical evolution of serology in 38 patients without evidence of hydatidosis

	Months after surgery					
	6	12	18	24	30	36
IgG +	100%	97.1%	90%	89.7%	88.9%	85.2%
IgM +	62.5%	25%	19%	12.5%	11.1%	8.6%
IgA +	52.5%	27.3%	21.7%	16.7%	11.5%	2.9%

TABLE 4. Post-surgical evolution of serology in 27 patients with cyst persistence

	Months after surgery					
	6	12	18	24	30	36
IgG +	100%	100%	100%	100%	100%	100%
IgM +	90.5%	81%	71.4%	61.9%	57.9%	42.9%
IgA +	87%	80%	84.2%	78.9%	72.2%	54.5%

TABLE 5. Post-chemotherapy evolution of serology in six patients with cyst persistence

	Months after surgery					
	6	12	18	24	30	36
IgG +	100%	100%	100%	100%	100%	100%
IgM +	75%	25%	25%	25%	25%	20%
IgA +	66.6%	66.6%	66.6%	66.6%	66.6%	20%

cysticercosis, leishmaniasis, toxoplasmosis, and trichinosis gave false-positive results with the different isotypes of immunoglobulins, with percentages between 8% (toxoplasmosis, leishmaniasis, and trichinosis) and 16% (cysticercosis), these values being lower than those reported by other authors. Orduña et al. (5), using crude and lipoprotein antigens, detected one false-positive result in two patients suffering cysticercosis. Matossian et al. (11) found 29% false-positives for ELISA-IgG and 7% for ELISA-IgM in patients with trichinosis, using crude hydatid fluid as antigen. The behaviour of the isotypes, depending on the state of the cyst, was similar to that described by other authors (3), and showed greater sensitivity in the complicated cysts.

All specific immunoglobulin isotypes from patients considered cured presented prolonged persistence, especially for the IgG. Although the IgA and IgM became negative most rapidly, it should be taken into account that their sensitivity in presurgical sera was insufficient, and that a negative trend was also found (although it was much slower) in patients with persistent cysts after

treatment. Such circumstances limit its usefulness in postsurgical monitoring.

The use of ELISA with enriched antigen fraction did not improve the posttreatment follow-up of the ELISA (22,23) or IHA (22,24,25) described to date. Nor did we find any relationship between the presence of IgM and the cyst status, in contrast to the results of Tassi et al. (1), who used a micro-ELISA with antigen fractions. However, in accordance with Tassi et al. (1), we related the postsurgical presence of this isotype to antigen release during surgery. Therefore, we conclude that following the serological evolution in each patient and verifying the persistence or decrease of specific antibody levels is the most effective serological alternative to treatment monitoring.

The data of the present study indicate that purification of the antigen improves specificity without diminishing sensitivity; however, no advantages are gained in the postsurgical follow-up of the patient.

REFERENCES

1. Tassi C, Baldelli F, Papili R, Valigi D. Specific IgM antibody response, evaluated by micro-ELISA, to various antigen fractions from sheep hydatid fluid and scolices of *E. granulosus*, in sera from patients affected by hydatid disease, before and after surgery. *Boll Ist Sieroter Milan* 1984;63:140-144.
2. Guisantes JA, Rubio M, Díaz R. Aplicación de un método inmunoenzimático (ELISA) al diagnóstico de la hidatidosis humana. *Bol Sanit Panam* 1981;90:160-168.
3. Navarrete N, Jersic MI, Denis R. Comparación de tres técnicas en el diagnóstico serológico de la hidatidosis humana. *Bol Chil Parasitol* 1995;50:97-100.
4. Njerhu FM, Gathuma JM, Okelo GBA, Tumboh-Oeri AG. Diagnosis of human hydatid disease in surgically-confirmed cases by use of the indirect haemagglutination test based on a thermo-stable lipoprotein and on unfractionated hydatid cyst fluid. *Am Trop Med Parasitol* 1989;83:299-303.
5. Orduña A, Castrodeza J, Gobernado C, Rodríguez-Torres A. Aplicación de un método inmunoenzimático al diagnóstico serológico de la hidatidosis humana. II. Especificidad y sensibilidad de las pruebas ELISA IgG e IgM específicas. *Laboratorio* 1985;476:113-126.
6. Schantz PM, Shanks D, Wilson M. Serologic cross-reactions with sera from patients with echinococcosis and cysticercosis. *Am J Trop Med Hyg* 1980;29:609-612.
7. Vicente F, Guisantes JA. Diagnóstico de la hidatidosis humana. Nuestra experiencia con cinco métodos inmunológicos. *Med Clin (Barc)* 1984;82:93-96.
8. Force L, Torres JM, Carrillo A, Buscá J. Evaluation of eight serological tests in the diagnosis of human echinococcosis and follow-up. *Clin Infect Dis* 1992;15:473-480.
9. Varela-Díaz VM, Coltorti EA, D'Alessandro A. Immunoelectrophoresis test showing *Echinococcus granulosus* arc 5 in human cases of *Echinococcus vogeli* and cysticercosis. *Am J Trop Med Hyg* 1978;27:554-557.
10. Verastegui M, Moro P, Guevara A, Rodríguez T, Miranda E, Gilman RH. Enzyme-linked immunoelectrotransfer blot test for diagnosis of human hydatid disease. *J Clin Microbiol* 1992;30:1557-1561.
11. Matossian RM, Radwan H, Meshefedjian GA. Immunoglobulin classes of reactive antibodies in diagnosis of human hydatidosis. *Helminth* 1991;28:41-47.
12. Bouchara JP, Robert P, Senet JM, Leynia P, Jarrige R. Mise en évidence de communautés antigéniques hôte-parasite dans l'hydatidose. *Bull Soc Path Exot* 1985;78:707-711.
13. Pezzella M, Galli C, Delia S, Solice F, Vullo V, Zennaro F. Fractionation and characterization of hydatid fluid antigens with identification of an antigen similar to human serum albumin. *Trans R Soc Trop Med Hyg* 1984;71:821-826.
14. Cameron GL, Satveley JM. Blood group P substance in hydatid cyst fluids. *Nature* 1957;179:147-148.
15. Makni S, Ayed KH, Dalix AM, Oriol R. Immunological localization of blood pl antigen in tissues of *Echinococcus granulosus*. *Ann Trop Med Parasitol* 1992;86:87-88.
16. Russi S, Siracusano A, Vicari G. Isolation and characterization of a blood PI active carbohydrate antigen of *Echinococcus granulosus* cyst membrane. *J Immunol* 1974;112:1061-1069.
17. Sbihi Y, Janssen D, Osuna A. Serologic recognition of hydatid cyst antigens using different purification methods. *Diagn Microbiol Infect Dis* 1996;24:205-211.
18. Sbihi Y, Janssen D, Osuna A. Specific recognition of hydatid cyst antigens by serum IgG, IgE, and IgA using western-blot. *J Clin Lab Anal* 1997;11:154-157.
19. Rogan MT, Craig PS, Zeyhle E, Romig T, Lubano GM, Deshan L. Evaluation of a rapid dot-ELISA as a field test for the diagnosis of cystic hydatid disease. *Trans R Soc Trop Med Hyg* 1991;85:773-777.
20. Oriol R, Williams JF, Pérez-Esandi MV, Oriol C. Purification of lipoprotein antigens of *Echinococcus granulosus* from sheep hydatid fluid. *Am J Trop Med Hyg* 1971;20:569-574.
21. Candolfi E, Kien T, Chaker E, Fourati M, Benyounes A. Intérêt de l'antigénémie et des anticorps IgG, IgM, IgA et IgE dans l'immunologie du kyste hydatique. Résultats de l'immunoenzymologie. *Bull Soc Path Ex* 1985;78:700-706.
22. Larrand L, Benito R, Sarría A, et al. Postoperative control for the persistence of positivity in three immunological tests used in the diagnosis of hydatidosis in children (E.L.I.S.A.-specific IgE, indirect hemagglutination and E.L.I.S.A.-total IgE). *Allergol Immunopathol* 1984;12:199-206.
23. Orduña A, Espinosa M, Bratos R, Ortiz de Lejárazu R, Rodríguez Torres A. Estudio de la evolución serológica postquirúrgica en pacientes con hidatidosis mediante pruebas clásicas y pruebas de ELISA. *Enferm Infecc Microbiol Clin* 1986;4:213-220.
24. Ravinder PT, Parija SC, Subba Rao KSVK. Evaluation of human hydatid disease before and after surgery and chemotherapy by demonstration of hydatid antigens and antibodies in serum. *J Med Microbiol* 1997;47:859-864.
25. Torres Rodríguez JM, Carrillo A, Casanovas M. Detección de anticuerpos IgE por un sistema ELISA de captura en la hidatidosis. *Med Clin (Barc)* 1989;92:37.